

Topological Borsuk problem

Yan Soibelman

February 1, 2008

To the memory of my wife Tanya with love

1 Introduction

This paper is based on my lecture at the US-Canada Mathcamp-2002 in Colorado Springs. The text is very elementary (it was designed for advanced high school students). At the same time the main problem about the topological Borsuk number for $\mathbf{R}^n, n > 2$ remains open since 1977.

I thank the organizers of the Mathcamp-2002 for inviting me to give lectures.

2 Borsuk conjecture and topological Borsuk number

Famous Borsuk conjecture says that any compact F in $\mathbf{R}^n$ can be partitioned into $n+1$ closed subsets of smaller diameter. The conjecture is true for $n = 2$ and $n = 3$ as well as for all F having smooth boundary. The latter result is topological. It is based on the following Borsuk theorem: if the standard sphere $S^{n-1} \subset \mathbf{R}^n$ is represented as a union $F_1 \cup F_2 \cup \dots \cup F_n$ of n closed subsets then at least one F_i contains a pair of antipodal points x and $-x$.

General Borsuk conjecture was disproved in [KK]. Let us call $b_{\mathbf{R}^n}(F)$ the minimal number of parts of smaller diameter necessary to partition F . Then Kahn and Kalai constructed such F that

$$b_{\mathbf{R}^n}(F) > (1.2)^{\sqrt{n}}.$$

The counterexample (as well as many of its simplifications) has combinatorial nature and deals with specific properties of the Euclidean metric in $\mathbf{R}^n$.

Topological version of the Borsuk problem was formulated around 1977 (see [So]). We recall it below.

Let (X, ρ_0) be a metric space (ρ_0 is the metric). For any compact $F \subset X$ we denote by $b_{(X, \rho_0)}(F) := b_X(F)$ its Borsuk number, i.e. the minimal number of parts of smaller diameter necessary to partition F . We denote by $B(X, \rho_0)$ the Borsuk number of the metric space X . By definition, any compact in X can be partitioned into $B(X, \rho_0)$ parts of smaller diameter, but there exist compacts which cannot be partitioned into $B(X, \rho_0) - 1$ such parts.

Let $\Omega(\rho_0)$ be the set of metrics on X which define the topology equivalent to the one given by ρ_0 . We will call elements of this set ρ_0 -equivalent metrics.

The number $B(X, \rho)$ can change as ρ varies inside of $\Omega(\rho_0)$.

Example 1 *Let ρ_0 be the standard Euclidean metric in $\mathbf{R}^2$ and ρ be the Minkowski metric, i.e. $\rho((x_1, y_1), (x_2, y_2)) = |x_1 - x_2| + |y_1 - y_2|$ in coordinates. Clearly $\rho \in \Omega(\rho_0)$. Then the classical result, which goes back to 1950's, says that $B(X, \rho) = 4$. On the other hand $B(X, \rho_0) = 3$.*

The following definition was given in [So].

Definition 1 *Topological Borsuk number of (X, ρ_0) is defined as*

$$B(X) = \min_{\rho \in \Omega(\rho_0)} B(X, \rho).$$

Topological Borsuk Problem. *Estimate $B(X)$ for the Euclidean space $X = \mathbf{R}^n$.*

In particular, is it true that $B(\mathbf{R}^n) \geq n + 1$? More speculatively, is it true that $B(\mathbf{R}^n)$ is bounded from below by $B(\mathbf{R}^n, \rho_0)$, where ρ_0 is the standard Euclidean metric?

3 2-dimensional case

In order to prove that $B(X) \geq m$ it suffices to prove that for any metric $\rho \in \Omega(\rho_0)$ there exists $c > 0$ and a finite subset $I \subset X$ consisting of m elements such that $\rho(i, j) = c$ for all $i \neq j$. If $X = \mathbf{R}^n$ and $\rho = \rho_0$ is the standard Euclidean metric then one can take $c = 1$, $m = n + 1$ and I be the

set of vertices of the regular simplex. It is not obvious that such I exists for other ρ_0 -equivalent metrics. In the case $n = 2$ the answer is positive due to the following theorem.

Theorem 1 ([So]) *Topological Borsuk number of the Euclidean $\mathbf{R}^2$ is equal to 3.*

The proof presented below is basically the same as in [So].

Proof. Let ρ be a metric on $X = \mathbf{R}^2$ which defines the topology equivalent to the standard one. Let us consider the map $f : X^3 \rightarrow \mathbf{R}^3$ such that

$$f(x_1, x_2, x_3) = (\rho_{12}, \rho_{23}, \rho_{13}),$$

where $\rho_{ij} = \rho(x_i, x_j), i \neq j$.

By our assumption the map f is continuous. It suffices to prove that the image of f intersects the line $l = \{(\rho_{12}, \rho_{23}, \rho_{13}) | \rho_{12} = \rho_{23} = \rho_{13}\}$ besides the obvious point $(0, 0, 0)$ which is the image of the diagonal $x_1 = x_2 = x_3$.

Notice that f is equivariant with respect to the natural actions of the cyclic group $\mathbf{Z}/3$ on X^3 and $\mathbf{R}^3$ (view $\mathbf{Z}/3$ as the subgroup of the symmetric group Σ_3 acting by permutations of coordinates).

It suffices to prove that there is no continuous $\mathbf{Z}/3$ -equivariant map between $X^3 \setminus f^{-1}(0)$ and $\mathbf{R}^3 \setminus l$.

Since $X^3 = \mathbf{R}^6$ and $f^{-1}(0) = \{(x_1, x_2, x_3) | x_1 = x_2 = x_3\}$ the set $X^3 \setminus f^{-1}(0)$ is equivariantly homotopic to the 3-dimensional sphere S^3 (which can be considered as a $\mathbf{Z}/3$ -invariant subset of the plane $x_1 + x_2 + x_3 = 0$). Similarly, $\mathbf{R}^3 \setminus l$ is equivariantly homotopic to the 1-dimensional sphere S^1 considered as a $\mathbf{Z}/3$ -invariant subset of the plane $\rho_{12} + \rho_{23} + \rho_{13} = 0$. Notice that the natural action of $\mathbf{Z}/3$ is free on both spheres. It is enough to prove that there is no $\mathbf{Z}/3$ -equivariant map between S^3 and S^1 . It was done in [So] by using the notion of the category (genus) of a topological space. Here we will use its modern version called the G -index of a topological space where G is a group acting on the space (see [M]). In our case $G = \mathbf{Z}/3$.

Definition 2 *Let G be a non-trivial finite group. An $E_n G$ -space is a G -space Y such that*

- a) G acts freely on Y ;*
- b) Y is n -dimensional;*
- c) Y is $(n - 1)$ -connected.*

If X is a G -space then $\text{ind}_G(X)$ (the G -index of X) is the minimal n such that there exists a G -equivariant map $X \rightarrow E_n G$ (the index can be infinite). It is easy to see that $\text{ind}_G(E_n G) = n$. One can prove (see [M], 6.2.5) that there is no G -equivariant continuous map $E_n G \rightarrow E_{n-1} G$ (Borsuk-Ulam type theorem). More generally, there is no G -equivariant continuous map $f : X \rightarrow Y$ between G -spaces X and Y such that $\text{ind}_G(X) > \text{ind}_G(Y)$.

On the other hand, if p is a prime number then any odd-dimensional sphere is a $\mathbf{Z}/p$ -space. Indeed, the group $\mathbf{Z}/p$ acts on $S^{2n-1} = \{(z_1, \dots, z_n) \in \mathbf{C}^n \mid \sum_i |z_i|^2 = 1\}$ via $(z_1, \dots, z_n) \mapsto (z_1 \exp(2\pi i/p), \dots, z_n \exp(2\pi i/p))$. Taking $p = 3$ we finish the proof of the theorem. ■

4 Conclusion

The proof above uses only elementary algebraic topology and very little information about the metric. The proof does not work when we replace $\mathbf{R}^2$ by $\mathbf{R}^n, n > 2$. In that case we have a Σ_{n+1} -equivariant continuous map $(\mathbf{R}^n)^{n+1} \rightarrow \mathbf{R}^{n(n-1)/2}$. Although arising topological spaces are still spheres, the natural actions of various subgroups of the symmetric group Σ_{n+1} are not free on the target. Perhaps one needs new ideas in order to estimate the topological Borsuk number of $\mathbf{R}^n$.

References

- [KK] J. Kahn, G. Kalai, A counterexample to Borsuk's conjecture. Bull. AMS, 29(1993), 60-62.
- [M] J. Matousek, Topological Methods in Combinatorics and Geometry (book, to appear).
- [So] Y. Soibelman, Calculation of a topological invariant connected with the partitioning of figures into parts of smaller diameter. (Russian) Mat. Zametki 27 (1980), no. 4, 647-649, 671. English Translation in Math. Notes 27 (1980), 3-4, 317-318.

Address: Department of Mathematics, KSU, Manhattan, KS 66506, USA
soibel@math.ksu.edu